

Package ‘WMWssp’

July 9, 2019

Type Package

Title Wilcoxon-Mann-Whitney Sample Size Planning

Version 0.4.0

Date 2019-07-09

Maintainer Martin Happ <martin.happ@aon.at>

Description Calculates the minimal sample size for the Wilcoxon-Mann-Whitney test that is needed for a given power and two sided type I error rate. The method works for metric data with and without ties, count data, ordered categorical data, and even dichotomous data. But data is needed for the reference group to generate synthetic data for the treatment group based on a relevant effect. For details, see Brunner, E., Bathke A. C. and Konietzschke, F: Rank- and Pseudo-Rank Procedures in Factorial Designs - Using R and SAS, Springer Verlag, to appear.

Depends R (>= 3.4.0)

License GPL-3

LazyData TRUE

URL <http://github.com/happma/WMWssp>

BugReports <http://github.com/happma/WMWssp/issues>

RoxygenNote 6.1.1

Suggests testthat

NeedsCompilation no

Author Arne C. Bathke [aut],
Edgar Brunner [aut],
Martin Happ [aut, cre] (<<https://orcid.org/0000-0003-0009-2665>>),
Frank Konietzschke [aut]

Repository CRAN

Date/Publication 2019-07-09 10:00:03 UTC

R topics documented:

WMWssp	2
WMWssp_maximize	3
WMWssp_minimize	4
WMWssp_noether	5

Index	7
--------------	----------

WMWssp	<i>Sample size calculation for the Wilcoxon-Mann-Whitney test.</i>
--------	--

Description

This function calculates the sample size for a given power, type-I error rate and allocation rate $t = n_1/N$. Additionally, the actual achieved power can be simulated.

Usage

```
WMWssp(x, y, alpha = 0.05, power = 0.8, t = 1/2,
       simulation = FALSE, nsim = 10^4)
```

Arguments

x	prior information for the first group
y	prior information for the second group
alpha	two sided type I error rate
power	power
t	proportion of subjects in the first group; or use t = "min" to use optimal proportion rate
simulation	TRUE if a power simulation should be carried out
nsim	number of simulations for the power simulation

Value

Returns an object from class WMWssp containing

result	A dataframe with the results.
t	The allocation rate which was used.
alpha	The type-I error rate which was used.
simulation	The achieved power in a simulation.
power	The power which was used.
N	The sample size needed.

References

Brunner, E., Bathke A. C. and Konietzschke, F. Rank- and Pseudo-Rank Procedures in Factorial Designs - Using R and SAS. Springer Verlag. to appear.

Happ, M., Bathke, A. C., & Brunner, E. (2019). Optimal Sample Size Planning for the Wilcoxon-Mann-Whitney-Test. *Statistics in medicine*, 38(3), 363-375.

Examples

```
# Prior information for the reference group
x <- c(315,375,356,374,412,418,445,403,431,410,391,475,379)
# generate data for treatment group based on a shift effect
y <- x - 20

# calculate sample size
ssp <- WMWssp(x, y, alpha = 0.05, power = 0.8, t = 1/2)
summary(ssp)
```

WMWssp_maximize	<i>Maximizing power for a given Type I error rate and prior information x and y</i>
-----------------	---

Description

This function maximizes the power of the Wilcoxon-Mann-Whitney test for a given total sample size N and type-I error rate with respect to the allocation rate $t = n_1/N$.

Usage

```
WMWssp_maximize(x, y, alpha = 0.05, N)
```

Arguments

x	a vector of prior information for the first group
y	a vector of prior information for the second group
alpha	Type I error rate
N	total sample size

Value

Returns an object from class WMWssp containing

result	A dataframe with the results.
t	The optimal allocation rate.
alpha	The type-I error rate which was used.
power	The maximized power.
N	The total sample size which was used.

References

Brunner, E., Bathke A. C. and Konietzschke, F. Rank- and Pseudo-Rank Procedures in Factorial Designs - Using R and SAS. Springer Verlag. to appear.

Happ, M., Bathke, A. C., & Brunner, E. (2019). Optimal Sample Size Planning for the Wilcoxon-Mann-Whitney-Test. *Statistics in medicine*, 38(3), 363-375.

Examples

```
# Prior information for the reference group
x <- c(315,375,356,374,412,418,445,403,431,410,391,475,379)
# generate data for treatment group based on a shift effect
y <- x - 20

#
N <- 112

# calculate optimal t
ssp <- WMWssp_maximize(x, y, alpha = 0.05, N)
summary(ssp)
```

WMWssp_minimize	<i>Minimizing samplesize for a given Type I and II error rate and prior information x and y</i>
-----------------	---

Description

This function minimizes the sample size for a given power and type-I error rate with respect to the allocation rate $t = n_1/N$.

Usage

```
WMWssp_minimize(x, y, alpha = 0.05, power = 0.8, simulation = FALSE,
  nsim = 10^4)
```

Arguments

x	a vector of prior information for the first group
y	a vector of prior information for the second group
alpha	Type I error rate
power	Power to detect a relative effect based on the prior information
simulation	TRUE if a power simulation should be carried out
nsim	number of simulations for the power simulation

Value

Returns an object from class WMWssp containing

result	A dataframe with the results.
t	The optimal allocation rate for minimizing the sample size.
alpha	The type-I error rate which was used.
power	The power which was used.
N	The minimized sample size.

References

Brunner, E., Bathke A. C. and Konietzschke, F. Rank- and Pseudo-Rank Procedures in Factorial Designs - Using R and SAS. Springer Verlag. to appear.

Happ, M., Bathke, A. C., & Brunner, E. (2019). Optimal Sample Size Planning for the Wilcoxon-Mann-Whitney-Test. *Statistics in medicine*, 38(3), 363-375.

Examples

```
# Prior information for the reference group
x <- c(315,375,356,374,412,418,445,403,431,410,391,475,379)
# generate data for treatment group based on a shift effect
y <- x - 20

# calculate optimal t
ssp <- WMWssp_minimize(x, y, alpha = 0.05, power = 0.8)
summary(ssp)
```

WMWssp_noether	<i>Sample size calculation for the Wilcoxon-Mann-Whitney test using the Noether formula</i>
----------------	---

Description

This function calculates the sample size for given type-I and type-II error probabilities using Noether's formula. If ties are present then prior information is needed.

Usage

```
WMWssp_noether(alpha, power, t, p, x = c(0), ties = FALSE)
```

Arguments

alpha	two sided type I error rate
power	power: detect a relative effect p at least with the specified power
t	proportion of subjects in the first group (between 0 and 1)
p	relative effect
x	prior information is only needed in case of ties
ties	TRUE if ties are possible (non continuous distribution), otherwise FALSE

Value

Returns an object from class WMWssp containing

result	A dataframe with the results.
t	The allocation rate which was used.
alpha	The type-I error rate which was used.
power	The power which was used.
N	The sample size needed.

References

Noether, G. E. (1987). Sample Size Determination for Some Common Nonparametric Tests. *Journal of the American Statistical Association* 85, 645.647.

Examples

```
# Prior information for the reference group
x <- c(315,375,356,374,412,418,445,403,431,410,391,475,379)
# generate data for treatment group based on a shift effect
y <- x - 20
# this data leads to a relative effect of p = 0.349

# calculate sample size for a balanced design
ssp <- WMWssp_noether(alpha = 0.05, power = 0.8, t = 1/2, p = 0.349)
summary(ssp)
```

Index

*Topic **export**

WMWssp, [2](#)

WMWssp_maximize, [3](#)

WMWssp_minimize, [4](#)

WMWssp_noether, [5](#)

WMWssp, [2](#)

WMWssp_maximize, [3](#)

WMWssp_minimize, [4](#)

WMWssp_noether, [5](#)