

Vignette for Risk and Performance Estimators Influence Functions Package

Anthony Christidis, R. Douglas Martin, Shengyu Zhang

May 14, 2021

Abstract

The RPEIF package computes the influence functions for risk and performance estimators based on the returns of individual assets or groups of assets, the latter for example in the context of an investment portfolio. The current version of the package treats five risk estimators and seven performance estimators, including among others, the standard deviation, value at risk, expected shortfall, Sharpe ratio, Sortino ratio estimators. The influence function of a risk or performance estimator provides a basis for computing an estimate of the finite-sample standard error when returns are serially uncorrelated, and also when the returns are serially correlated, as discussed in [Chen and Martin, 2021]. This RPEIF package is used in the package RPESE to compute the time series of influence function transformed returns needed to compute the standard errors of risk and performance estimators

1 Influence Functions Theoretical Background

The Influence Functions package (RPEIF) makes use of the definition and theory of influence functions in robust statistics, as introduced by [Hampel, 1974], further treated in [Hampel et al., 1986]. Recent use of influence functions for the analysis of parametric and nonparametric expected shortfall (ES) is discussed in [Martin and Zhang, 2019], and use of influence functions for computing standard errors of risk and performance estimators with serially correlated returns is discussed in [Chen and Martin, 2021]. Here we provide the definition and basic properties of influence functions, with a view toward their use for understanding the influence of outliers on risk and performance estimators, and for computing standard errors of such estimators for both uncorrelated and serially correlated returns.

1.1 Risk and Performance Estimator Functional Representations

The large-sample value (as sample size n tends to infinity) of a risk or performance estimator may be represented as a functional $T = T(F)$ of the marginal distribution function F of a time series r_1, r_2, \dots, r_n of returns.¹ For example the functional for the mean (expected value) is

¹The term functional refers to a function whose domain is an infinite dimensional space, e.g., the space of distribution functions.

$$\mu(F) = \int r dF(r) \quad (1)$$

and the functional for the standard deviation (returns volatility) is

$$\sigma(F) = \left[\int (r - \mu(F))^2 dF(r) \right]^{\frac{1}{2}}. \quad (2)$$

Given a functional representation $T(F)$ of an estimator, a finite-sample *non-parametric* estimator T_n is easily obtained by replacing the unknown distribution F by the empirical distribution F_n that has a jump of height $1/n$ at each of the observed returns values r_1, r_2, \dots, r_n :

$$T_n = T(F_n) = T(r_1, r_2, \dots, r_n). \quad (3)$$

For example, the finite-sample non-parametric estimators of the mean and standard deviation are the sample mean and sample standard deviation, respectively:

$$\hat{\mu}_n = \frac{1}{n} \sum_{t=1}^n r_t \quad \hat{\sigma}_n = \left[\frac{1}{n} \sum_{t=1}^n (r_t - \hat{\mu}_n)^2 \right]^{\frac{1}{2}}. \quad (4)$$

We note that one can also derive parametric estimators from parametric functional representation obtained by replacing F by F_θ , where θ is the parameter vector for a parametric distribution function. In this case one obtains the finite-sample estimator by replacing the unknown parameter by its estimator, typically the maximum-likelihood estimator (MLE). See for example, [Martin and Zhang, 2019] for a treatment of parametric and non-parametric ES estimators for normal and t-distributions. However, this first version of the RPEIF package only deals with non-parametric risk and performance estimators.

1.2 Estimator Influence Function Definition

Influence Functions are based on the use of the following mixture distribution perturbation of a fixed target distribution $F(x)$:

$$F_\gamma(x) = (1 - \gamma)F(x) + \gamma\delta_r(x), \quad 0 \leq \gamma < 1/2 \quad (5)$$

where $\delta_r(x)$ is a point mass discrete distribution function with a jump of height one located at value r . The influence function of an estimator with functional form $T(F)$ is defined as:

$$IF(r; T, F) = \lim_{\gamma \rightarrow 0} \frac{T(F_\gamma) - T(F)}{\gamma} = \frac{d}{d\gamma} T(F_\gamma)|_{\gamma=0} \quad (6)$$

The influence function is a special directional derivative (i.e., a Gateaux derivative) of the functional $T(F)$ in the direction of a point mass distributions δ_r , evaluated at F .

It is straightforward, and more or less tedious, to derive formulas for the influence functions of risk and performance estimators. For example, the influence function of the sample mean is:

$$IF(r; \mu; F) = r - \mu \quad (7)$$

where $\mu = \mu(F)$ depends on the underlying returns marginal distributon F . The above influence function has the property that its expected value is zero, which is a reflection of the general property than an influence function has zero expected value:

$$E [IF(r; T, F)] = 0. \quad (8)$$

1.3 A Key Influence Function Property

A key influence function property is that for well behaved estimator functionals, the finite-sample estimator $T_n = T(F_n) = T(r_1, r_2, \dots, r_n)$ can be expressed as the following linear combination of influence-function transformed returns:

$$T_n - T(F) = \frac{1}{n} \sum_{t=1}^n IF(r_t; T, F) + remainder \quad (9)$$

where the $IF(r_t; T, F)$ are influence-function (IF) transformed returns, and the remainder goes to zero in a probabilistic sense as $n \rightarrow \infty$. Thus the finite sample variance of T_n is approximately given by

$$Var(T_n) = Var \left[\frac{1}{n} \sum_{t=1}^n IF(r_t; T, F) \right] \quad (10)$$

and in the special case where the returns r_t are i.i.d., the IF -transformed returns are i.i.d., and the variance of T_n reduces to

$$Var(T_n) = \frac{1}{n} E[IF^2(r_1; T, F)]. \quad (11)$$

and the expectation on the right-hand side can be evaluated empirically as the sample mean of the $IF(r_t; T, F)$, $t = 1, 2, \dots, n$.

However, when the r_t , $t = 1, 2, \dots, n$ are serially correlated, the IF -transformed returns time series $IF(r_t; T, F)$ will generally have serial correlation that needs to be accounted for in calculating the variance on the the right-hand-side of (10). Spectral analysis theory, extensively used in science and engineering, shows that the variance of the sum of the values of a serially correlated stationary time series is given by the spectral density of the time series at zero frequency. Thus the problem of estimating the variance (10), with possibly serially correlated returns, reduces to the problem of estimating the spectral density at zero frequency at zero frequency of the the time series $IF(r_t; T, F)/n$. [Chen and Martin, 2021] show how to do this by a polynomial gneralized linear model (GLM) fitting method, with elastic net (EN) regularization, that works well when the returns are serially correlated, as well as when they are uncorrelated. Their methodology, which we refer to as the CM method, is implemented in the Risk and Performance Estimator Standard Error (RPESE) package, which in turn makes fundamental use of the RPEIF package we are discussing here.

2 RPEIF Package Estimators and their Influence Functions

Table 1 lists the symbolic or acronym names and descriptions of the risk and performance estimators for which the RPEIF package computes influence functions. Each estimator has a functional representation, a sample based estimator as a function of a time series of asset returns, and an influence function formula. We illustrate this below for the case of the standard deviation and Sharpe ratio estimators.

| Name | Estimator Description |
|-------------|---|
| μ | Mean |
| μ_M | M-estimator of location with efficiency e |
| σ | Standard deviation |
| SSD | Semi-standard deviation |
| LPM | Lower partial moment of order 1 or 2 with threshold c |
| ES | Expected shortfall with tail probability α |
| VaR | Value-at-risk with tail probability α |
| SR | Sharpe ratio |
| DSR | Downside Sharpe ratio |
| SoR | Sortino ratio with threshold the mean or a constant c |
| ESratio | Mean excess return to ES ratio with tail probability α |
| VaRratio | Mean excess return to VaR ratio with tail probability α |
| OmegaRatio | Omega ratio with threshold c |
| RachevRatio | Rachev ratio with lower upper tail probabilities α and β |

Table 1: Estimator Names and Descriptions

Standard Deviation (SD)

The functional representation $\sigma(F)$ of the sample standard deviation estimator is given by (2), and the formula for the influence function of the sample standard deviation estimator is

$$IF(r; \sigma; F) = (2\sigma)^{-1}((r - \mu)^2 - \sigma^2) \quad (12)$$

where $\mu = \mu(F)$ and $\sigma = \sigma(F)$. The sample standard deviation estimator is given in (4).

Sharpe Ratio (SR)

The formula for the functional representation of the Sharpe ratio is

$$SR(F) = \frac{\mu(F) - r_f}{\sigma(F)} = \frac{\mu_e(F)}{\sigma(F)} \quad (13)$$

where $\mu_e(F)$ is the mean excess return over the risk-free rate r_f , and the SR influence function formula is:

$$IF(r; SR; F) = -\frac{\mu_e}{2\sigma^3}(r - \mu)^2 + \frac{1}{\sigma}(r - \mu) + \frac{\mu_e}{2\sigma}. \quad (14)$$

The sample estimator of the Sharpe ratio is:

$$SR_n = \frac{\hat{\mu}_n - r_f}{\hat{\sigma}_n} \quad (15)$$

where $\hat{\mu}_n$ and $\hat{\sigma}_n$ are the sample mean and sample standard deviation.

The functional forms, and derivations of the influence function formulas for all the estimators in Table 1 may be found in Zhang et al. [2020]. For the convenience of interested readers, the influence functions of all the estimators in Table 1 are provided in Appendix A.

Nuisance Parameters

We note that the influence function formulas above, as well as in Appendix A, contain one or more *nuisance* parameters that need to be specified in order to compute influence function values for various values of a return r . For example, the IF of the sample standard deviation depends on the two nuisance parameters $\mu = \mu(F)$ and $\sigma = \sigma(F)$, and the Sharpe ratio IF depends on the three nuisance parameters $\mu = \mu(F)$, $\sigma = \sigma(F)$, and $SR = SR(F)$. The third column of Table 2 contains the nuisance parameters for each risk and performance estimator, along with the user specified parameters such as tail probabilities, Sortino ratio threshold, etc. The fourth column of Table (2) contains the values of the nuisance parameters for the case of monthly returns distribution with mean $\mu = 0.01$, volatility $\sigma = 0.05$ and risk-free rate $r_f = 0$. The formulas for the nuisance parameters and R code to calculate their values are contained in Appendix B.

3 Using the RPEIF Package to Evaluate Influence Functions and Compute Influence-Function Transformed Returns

Load and install and load the RPEIF package with the code:

```
library(devtools)
install_github("AnthonyChristidis/RPEIF")
```

The RPEIF package computes the influence functions time series of the returns for the risk and performance measures in RPESE, which in turn is used in PerformanceAnalytics if the user wishes to return the standard errors for the risk and performance measures. The RPEIF package is therefore required by the RPESE package.

To demonstrate the usage of the RPEIF package, we will make use of the `edhec` data set available from `PerformanceAnalytics`. This data set contains hedge fund returns from January, 1997 to August, 2008. You can load that data set, and confirm that it is an `xts` time series object with the following R code.

```
library(RPEIF)
data(edhec, package = "PerformanceAnalytics")
class(edhec)

## [1] "xts" "zoo"
```

Use of the code line

```
colnames(edhec)
```

will reveal to you that the hedge fund style names are too long to display well in plots. So use the following code to replace those long names with shorter names as follows, and view the first six values of the CA fund.

```
colnames(edhec) <- c("CA", "CTAG", "DIS", "EM", "EMN", "ED", "FIA", "GM", "LS", "MA",
  "RV", "SS", "FoF")
colnames(edhec)

## [1] "CA" "CTAG" "DIS" "EM" "EMN" "ED" "FIA" "GM" "LS"
## [10] "MA" "RV" "SS" "FoF"

head(edhec[, "CA"])

## [1] 0.0119 0.0123 0.0078 0.0086 0.0156 0.0212
```

To see what functions are contained in the RPEIF package, use the code line:

```
library(RPEIF)
ls("package:RPEIF")
```

The list of functions you see as a result of the above code will include the functions shown in the second column of [2](#). We use the term `IF.xxxx` as a generic for any one of the functions in the second column. The third column lists the names of nuisance parameters that need to be specified one way or another to compute the values of an influence function at various values of the influence function argument. The fourth column contains normal distribution “typical” values for the nuisance parameters.

You can get help on any of the functions `IF.xxxx` with a command like the following:

```
# Help file for the IF function
help(IF.SD)
```

| Estimator | IF Function | Nuisance Parameters | Normal Dist. Nuisance Pars |
|---------------------------------|----------------|---|--------------------------------------|
| μ | IF.Mean | μ | .01 |
| μ_M | IF.robMean | $\mu, \sigma, E(\psi')$ | .01, .05, .9038 |
| σ | IF.SD | μ, σ | .01, .05 |
| SSD | IF.SemiSD | $\mu, SSD, SMEAN$ | .01, .0354, -0.0199 |
| LPM1 | IF.LPM | $LPM1_c$ | .0153 |
| LPM2 | IF.LPM | $LPM2_c$ | .0009 |
| ES | IF.ES | q_α, ES_α | -.0541, .0777 |
| VaR | IF.VaR | $q_\alpha, f(q_\alpha)$ | -.0541, 3.5100 |
| SR | IF.SR | $\mu_e, \sigma, SR,$ | .01, .05, .20 |
| DSR | IF.DSR | $\mu, SSD, SMEAN, DSR$ | .01, .0354, -.0199, .2828 |
| SoR _c | IF.SoR | $\mu, LPM2_c, SoR_c$ | .01, .0009, .3336 |
| SoR _{μ} | IF.SoR | $\mu, SSD, SMEAN, SoR_\mu$ | .01, .0354, -.0199, .2828 |
| ESratio | IF.ESratio | $\mu, q_\alpha, ES_\alpha, ESratio$ | .01, -.0541, .0777, .1286 |
| VaRratio | IF.VaRratio | $\mu, q_\alpha, f(q_\alpha), VaRratio$ | .01, -.0541, 3.5100, .1849 |
| RachevRatio | IF.RachevRatio | $q_\alpha, ES_\alpha, q_{1-\beta}, EG_\beta, RachR$ | -.0541, .0777, .0741, 0.0977, 1.2572 |
| OmegaRatio | IF.OmegaRatio | $LPM1_c, UPM1_c, \Omega$ | .0153, .0253, 1.6516 |

Table 2: IF R Code Functions and Corresponding Nuisance Parameters

The R `IF.xxxx` functions are used for two distinct purposes. The first is to evaluate an estimator influence function at a set of argument values, and plot them to display the shape of the influence function. This allows the user to explore the different shapes of the influence functions of different estimators. The second and primary purpose of the functions is to compute influence-function transformed time series of returns, as a first step in the overall method of computing standard errors for risk and performance estimators.

The arguments of the R functions in Table (2) are all the same, except for the nuisance parameters which are estimator specific. For example, the arguments of `IF.SD` and `IF.SR` are as follows.

```
args(IF.SD)

## function (returns = NULL, evalShape = FALSE, retVals = NULL,
##      nuisPars = NULL, k = 4, IFplot = FALSE, IFprint = TRUE, prewhiten = FALSE,
##      ar.prewhiten.order = 1, cleanOutliers = FALSE, cleanMethod = c("locScaleRob")[1],
##      eff = 0.99, ...)
## NULL

args(IF.SR)
```

```
## function (returns = NULL, evalShape = FALSE, retVals = NULL,
##      nuisPars = NULL, k = 4, IFplot = FALSE, IFprint = TRUE, rf = 0,
##      prewhiten = FALSE, ar.prewhiten.order = 1, cleanOutliers = FALSE,
##      cleanMethod = c("locScaleRob")[1], eff = 0.99, ...)
## NULL
```

You see that the arguments of the two functions are the same except for the nuisance parameters arguments `parsSD`.`IF` and `parsSR`.`IF`.

Note that you get no results using an `IF.xxxx` function with no arguments, for example try `IF.Mean()` and see what you get. You need to either set `evalShape=T`, or supply a returns xts object for the argument `returns`, in order for an `IF.xxxx` function to compute results.

3.1 Evaluating and Plotting Influence Functions for Shape Comparisons

In order to compute the values and plot the shapes of influence functions using the R `IF.xxxx` functions in Table 2, the values of the nuisance parameters in the third column of 2 need to be specified, and there are two basic methods of doing so. The first method is to use “typical” values for those parameters in the case of a specific returns distribution. Such values are provided for the case of a normal returns distribution in the fourth column of the Table 2. The second method is to provide a time series of returns of an asset of interest as one of the function’s arguments, and in that case the R function will estimate the values of the nuisance parameters from the time series of returns.

Evaluating and Plotting Influence Functions Using Default Nuisance Parameter Values

Using the argument `evalShape=T` and `IFplot=T` as follows, you can easily get influence function plots. For example, you get plots of the influence functions of the standard deviation and Sharpe ratio estimators using the following code.

```
par(mfrow = c(1, 2), pty = "s")
outSD <- IF.SD(evalShape = T, IFplot = T, IFprint = T)
outSR <- IF.SR(evalShape = T, IFplot = T, IFprint = T)
par(mfrow = c(1, 1))
```

The plots are shown in Figure 1.




Figure 1: Influence Functions of Standard Deviation and Sharpe Ratio (“typical” nuisance parameters values)

Evaluating and Plotting Influence Functions with User-Specified Nuisance Parameter Values

Using the `nuisParsFn` function, you can generate nuisance parameters by specifying “typical” values based on some assumed returns distribution. For the risk measure estimators and performance measure estimators, the `nuisParsFn` function assumes follow the normal distribution, with monthly mean return of 1% and risk-free rate $r_f = 0$, and monthly volatility of 5% (the corresponding annual mean and volatility are 12% and 17.3%, respectively). We also note that we use $\alpha = 0.10$ for quantiles, VaR and ES, use that value along with the value $\beta = 0.10$ for the Rachev ratio, and use $c = 0$ for lower partial moments and Sortino ratio with fixed threshold.

To see the default values from the `nuisParsFn` function, you can look at the arguments of the function. The default arguments are the ones used to generate the values in Table 2.

```
args(nuisParsFn)

## function (mu = 0.01, sd = 0.05, c = 0, alpha = 0.1, beta = 0.1)
## NULL
```

To generate nuisance parameters using different default settings for any of the arguments, you can specify the new values in the arguments of the function. For example, if you want to generate nuisance parameters by using a mean return of 2% instead of 1% and a volatility of 15% instead of 5% (the defaults) to generate the influence functions plots for the standard deviation and the Sharpe ratio, you can use the `nuisPars` argument as in the following code.

```

par(mfrow = c(1, 2), pty = "s")
outSD <- IF.SD(evalShape = T, IFplot = T, IFprint = T, nuisPars = nuisParsFn(mu = 0.02,
  sd = 0.15))
outSR <- IF.SR(evalShape = T, IFplot = T, IFprint = T, nuisPars = nuisParsFn(mu = 0.02,
  sd = 0.15))
par(mfrow = c(1, 1))

```




Figure 2: Influence Functions of Standard Deviation and Sharpe Ratio (user-specified nuisance parameters)

Evaluating and Plotting Influence Functions Using Nuisance Parameter Values Estimated from a Returns Time Series

You can also compute influence function shapes using nuisance parameter values that are estimated from a returns time series of interest, the latter of which is specified by using the argument `returns` of the `IF.xxxx` functions. For example, you could use the CA hedge fund time series for this purpose by using the following code, the results of which are shown in Figure 3. Note that in using the `IF.SD` and `IF.SR` functions, the `evalShape=T` is retained.

```

library(xts) # Must have installed xts package
retCA <- edhec$CA
par(mfrow = c(1, 2), pty = "s")
outSD <- IF.SD(returns = retCA, evalShape = T, IFplot = T, IFprint = T)
outSR <- IF.SR(returns = retCA, evalShape = T, IFplot = T, IFprint = T)
par(mfrow = c(1, 1))

```




Figure 3: Influence Functions of Standard Deviation and Sharpe Ratio (estimated nuisance parameters values)

3.2 Evaluating and Plotting Influence-Function Transformed Returns Time Series

The general method of estimating standard errors of risk and performance estimators, described in [Chen and Martin, 2021], consists of computing a good estimate of the variance of time series sample average of influence-function transformed returns shown on the right-hand-side of (10). In case you are interested in seeing what influence-function transformed returns look like for actual returns data, can easily compute such returns with the `IF.xxxx` function. For example, you can do so for SD and SR estimators for the CA hedge fund returns with the following code, whose results for the CA returns, and for the SD and SR estimator influence function transformed returns are shown in Figure 4.

```
retCA <- edhec$CA
outSD <- IF.SD(returns = retCA, IFplot = T, IFprint = T)
outSR <- IF.SR(returns = retCA, IFplot = T, IFprint = T)
par(mfrow = c(3, 1))
plot(retCA, lwd = 0.8, ylab = "Returns", main = "CA Hedge Fund Returns")
plot(outSD, lwd = 0.8, main = "IF.SD Transformed Returns")
plot(outSR, lwd = 0.8, main = "IF.SR Transformed Returns")
```




Figure 4: Standard Deviation and Sharpe Ratio Influence-Function Transformed Returns for CA Hedge Fund

4 Outlier Cleaning

A very reliable outlier cleaning method that shrinks outliers can be obtained based on a robust location M-estimator and an associated robust scale estimator \hat{s} . A location M-estimator is computed as a solution of the equation

$$\sum_{t=1}^n \psi \left(\frac{r_t - \hat{\mu}_M}{\hat{s}} \right) = 0 \quad (16)$$

where $\psi = \psi(x)$ is a suitable odd “psi” function, and \hat{s} is a robust scale estimate of the residuals $\epsilon_t = r_t - \hat{\mu}_M$. For an introduction to location M-estimators and their computation, see Sections 2.3 and 2.7 of [Maronna et al., 2019].

Here we use the special optimal psi function

$$\psi_{opt}(x) = SGN(x) \left(|x| + \frac{a}{\phi(x)} \right)^+, \text{ with } a = 0.002449 \quad (17)$$

where $\phi(x)$ is the standard normal density function, and a plot of this psi function is displayed in Figure 5. The function $\psi_{opt}(x)$ has a “smooth rejection” character in that the function is continuous, and all returns with a scaled residual $(r_t - \hat{\mu}_M)/\hat{s}$ larger in magnitude than 3.568 are “rejected”, in the sense that such returns have no influence on the estimate $\hat{\mu}_M$ since such terms are set to zero in (16). For further details on the function $\psi_{opt}(x)$, see Section 5.8 of [Maronna et al., 2019].




Figure 5: Optimal Bias Robust Psi Function with $\psi_{opt}(x) = 0$ for $|x| > 3.568$ and 99% Normal Distribution Efficiency.

Based on a location estimate $\hat{\mu}_M$ and associated scale estimate \hat{s} , it is natural to define returns outliers as those returns r_t that fall outside the interval $[\hat{\mu}_M - 3.568 \cdot \hat{s}, \hat{\mu}_M + 3.568 \cdot \hat{s}]$. Such outliers are then “cleaned” by shrinking them to the nearest boundary of that interval.

For the fixed income arbitrage (FIA) hedge fund returns, it turns out that $\hat{\mu}_M = .00640$ and $\hat{s} = .00547$ is $[-.0131, .0259]$. Correspondingly, FIA returns with values less than $-.0131$, or greater than $.0259$, are detected as outliers and shrunk accordingly. The following code results in Figure 6, which shows the sample mean IF transformed FIA returns (which are equal to FIA returns minus the very small mean of the FIA returns) in the top plot, and the outlier cleaned IF tranformaed FIA returns in the bottom plot.

```
retFIA <- edhec$FIA
iftrFIA <- IF.Mean(returns = retFIA, IFprint = T)
iftrFIAClean <- IF.Mean(returns = retFIA, cleanOutliers = T, eff = 0.99, IFprint = T)
par(mfrow = c(2, 1))
plot(iftrFIA, main = "FIA Returns", lwd = 0.8)
plot(iftrFIAClean, main = "Outlier Cleaned FIA Returns", lwd = 0.8)
par(mfrow = c(1, 1))
```




Figure 6: IF and Outlier Cleaned IF Transformed FIA Returns

Just as a check to compare the IF transformed and outlier cleaned IF transformed FIA returns, we show in Figure 7 the former plot with a solid line, and the latter as overlaid dots.

```
plot(iftrFIA, lwd = 0.8, main = "FIA Returns (line) and Outlier Cleaned FIA Returns (dots)")
points(iftrFIAClean, pch = 16)
```




Figure 7: Overlaid IF (line) and Outlier Cleaned IF (dots) Transformed FIA Returns

5 Robust Estimator of Mean Influence Function

As an alternative to the outlier cleaning method described in Section 4, the influence function for a robust M-estimator of location (mean) is available in RPEIF. A rigorous treatment of the influence function of location M-estimators can be found in [Maronna et al., 2019]. In particular, the influence function of M-estimators of location μ_M is given by

$$IF(r; \mu_M; F) = \sigma \frac{\psi((r - \hat{\mu}_\infty)/\sigma)}{E_F[\psi'((r - \hat{\mu}_\infty)/\sigma)]} \quad (18)$$

where the asymptotic value $\hat{\mu}_\infty$ satisfies

$$E_F((r - \hat{\mu}_\infty)/\sigma) = 0, \quad (19)$$

and σ is a known dispersion parameter. For a previously computed dispersion estimator $\hat{\sigma}$, if F is symmetric, then the IF simplifies to

$$IF(r; \mu_M; F) = \hat{\sigma}_\infty \frac{\psi((r - \hat{\mu}_\infty)/\hat{\sigma}_\infty)}{E_F[\psi'((r - \hat{\mu}_\infty)/\hat{\sigma}_\infty)]}. \quad (20)$$

Replacing $\hat{\mu}$ and $\hat{\sigma}_\infty$ by their robust sample estimates $\hat{\mu}_M$ and \hat{s}_{mad} , (20) can thus be formulated for a finite sample as

$$IF(r_t; \hat{\mu}_M; F_n) = \hat{s}_{mad} \frac{\psi((r_t - \hat{\mu}_M)/\hat{s}_{mad})}{\frac{1}{n} \sum_{t=1}^n [\psi'((r_t - \hat{\mu}_M)/\hat{s}_{mad})]}, \quad (21)$$

where

$$\hat{s}_{mad} = MAD(\mathbf{r}) = 1.4826 \cdot MED\{|r_t - MED(\mathbf{r})|\}_{t=1}^n. \quad (22)$$

In RPEIF, the families of ψ functions are the same ones available in the package `RobStatTM`, namely `mOpt`, `opt` and `bisquare`. The default is the `mOpt` family, where

$$\psi_{mOpt}(x) = \begin{cases} x & |x| \leq 1 \\ \frac{\phi(1)}{\phi(1)-a} \left(x - SGN(x) \frac{a}{\phi(x)} \right) U(c - |x|) & |x| > 1 \end{cases} \quad (23)$$

where $\phi(x)$ is the standard normal density function, $U(x)$ is the unit step function, and the constants a and c depend on the desired normal distribution efficiency. These constants are computed internally in RPEIF for a desired efficiency.

The following code computes, and plots in Figure 8, the IF.Mean transformed FIA returns, and the IF.robMean transformed FIA returns.

```
retFIA <- edhec$FIA
iftrFIA <- IF.Mean(returns = retFIA, IFprint = T)
iftrFIARob <- IF.robMean(returns = retFIA, family = "mopt", eff = 0.95, IFprint = T)
par(mfrow = c(2, 1))
plot(iftrFIA, main = "IF.Mean Transformed FIA Returns", lwd = 0.8)
plot(iftrFIARob, main = "IF.robMean Transformed FIA Returns", lwd = 0.8)
par(mfrow = c(1, 1))
```





Figure 8: IF.Mean and IF.robMean FIA Transformed FIA Returns

Just as a check to compare the non-robust and robust IF tranformed FIA returns, we show in Figure 9 the former plot with a solid line, and the latter as overlaid dots.

```
plot(iftrFIA, lwd = 0.8, main = "FIA Returns (line) and Outlier Cleaned FIA Returns (dots)")
points(iftrFIARob, pch = 16)
```




Figure 9: IF.Mean (line) and IF.robMean (dots) Transformed FIA Returns

6 Prewhitening

Spectral density function estimation is a frequently used method in the field of signal processing, and in other engineering and science applications. Prewhitening is a technique often used to improve the performance of spectral density estimators. Since the core of the method described in [Chen and Martin, 2021] is estimation of a spectral density at frequency zero of an influence-function transformed returns time series IF_t , it is natural to be able to use prewhitening of that time series. The `seCorIFPW` variant of the basic `seCorIF` method in the `RPESE` package implement such prewhitening.

A prewhitened version IF_t^{pw} of the IF_t time series is computed as

$$IF_t^{pw} = IF_t - \hat{\rho}IF_{t-1} \quad (24)$$

where $\hat{\rho}$ is a lag-one serial correlation coefficient estimat of the IF_t . In general the IF_t^{pw} series is not a serially uncorrelated (white noise) series, but it has considerably less serial correlation than IF_t , and a periodogram estimator based on IF_t^{pw} will suffer from relatively little bias compared with one based on IF_t .

The following code computes, and plots in Figure 10, the IF.SR transformed FIA returns, and the prewhitened IF.SR transformed FIA returns.

```

iftrFIA <- IF.SR(returns = retFIA, IFprint = T)
PWiftrFIA <- IF.SR(returns = retFIA, prewhiten = T, IFprint = T)
par(mfrow = c(2, 1))
plot(iftrFIA, main = "IF.SR Transformed FIA Returns", lwd = 0.8)
plot(PWiftrFIA, main = "Prewhitened IF.SR Transformed FIA Returns", lwd = 0.8)
par(mfrow = c(1, 1))

```




Figure 10: IF.SR Transformed FIA Returns (top) and Prewhitened Version of the Same (bottom)

The following code computes and displays the lag-1 through lag-5 autocorrelations of the time series in Figure 10:

```

corr.iftrFIA <- acf(iftrFIA, lag.max = 5, plot = F)
corr.PWiftrFIA <- acf(PWiftrFIA, lag.max = 5, plot = F)
out <- data.frame(round(rbind(corr.iftrFIA$acf, corr.PWiftrFIA$acf), 2))
names(out) <- c("Lag0", "Lag1", "Lag2", "Lag3", "Lag4", "Lag5")
row.names(out) <- c("autocorr: IF.SR FIA ret", "autocorr: PW IF.SR FIA ret")
out[-1]

##                Lag1 Lag2 Lag3 Lag4 Lag5

```

```
## autocorr: IF.SR FIA ret    0.44  0.10  0.01  0.01 -0.06
## autocorr: PW IF.SR FIA ret 0.05 -0.09 -0.05  0.04 -0.07
```

Outlier Cleaning Prior to Prewhitening

Since outliers can have adverse influence not only on risk and performance estimators, but also on the estimator $\hat{\rho}$ used for prewhitening, it is always a good idea to apply the outlier cleaning method of Section 4 before prewhitening. We illustrate doing this with the following code and plots, and autocorrelation calculations.

```
iftrFIAcl <- IF.Mean(returns = retFIA, cleanOutliers = T, IFprint = T)
PWiftrFIAcl <- IF.Mean(returns = retFIA, cleanOutliers = T, prewhiten = T, IFprint = T)
par(mfrow = c(2, 1))
plot(iftrFIAcl, type = "b", main = "FIA Outlier Cleaned Returns", lwd = 0.8)
plot(PWiftrFIAcl, type = "b", main = "Prewhitened FIA Outlier Cleaned Returns", lwd = 0.8)
par(mfrow = c(1, 1))
```




Figure 11: IF.SR Transformed Outlier Cleaned FIA Returns (top) and Prewhitened Version of the Same (bottom)

The following code computes and displays the lag-1 through lag-5 autocorrelations of the time series in Figure 11:

```
corr.iftrFIAcl <- acf(iftrFIAcl, lag.max = 5, plot = F)
corr.PWiftrFIAcl <- acf(PWiftrFIAcl, lag.max = 5, plot = F)
out <- data.frame(round(rbind(corr.iftrFIAcl$acf, corr.PWiftrFIAcl$acf), 2))
names(out) <- c("Lag0", "Lag1", "Lag2", "Lag3", "Lag4", "Lag5")
row.names(out) <- c("autocorr: IF.SR Cleaned FIA ret", "autocorr: PW IF.SR Cleaned FIA ret")
out[, -1]

##                               Lag1 Lag2 Lag3 Lag4 Lag5
## autocorr: IF.SR Cleaned FIA ret   0.41  0.19  0.26  0.14  0.00
## autocorr: PW IF.SR Cleaned FIA ret -0.02 -0.06  0.20  0.07 -0.06
```

Appendix A. Estimator Functional Forms and Influence Function Formulas

Here we provide the functional form of all the estimators in Table 2, and the formulas for their influence functions. The interested reader can find the derivations of these formulas in [Zhang, 2009].

Mean

The functional representation $\mu(F)$ of the sample mean estimator is given by (1), and the formula for the influence function of the sample mean is:

$$IF(r; \mu; F) = r - \mu \tag{25}$$

where $\mu = \mu(F)$.

Standard Deviation

The functional form of standard deviation is

$$\sigma(F) = \left(\int (x - \mu(F))^2 dF(x) \right)^{1/2}. \tag{26}$$

and the formula for the influence function of the standard deviation is:

$$IF(r; \sigma; F) = \frac{1}{2\sigma} \left((r - \mu)^2 - \sigma^2 \right) \tag{27}$$

Semi-Standard Deviation (SemiSD)

The functional representation $SSD(F)$ of the sample semi-standard deviation is

$$SSD(F) = \left(\int_{-\infty}^{\mu(F)} (x - \mu(F))^2 dF(x) \right)^{1/2} \quad (28)$$

and the formula for the influence function of the sample semi-standard deviation is:

$$IF(r; SSD; F) = \frac{(r - \mu)^2 \cdot \mathbf{I}(r \leq \mu) - 2 \cdot SMEAN \cdot (r - \mu) - SSD^2}{2 \cdot SSD} \quad (29)$$

where $\mu = \mu(F)$, $SSD = SSD(F)$, and $SMEAN$ is the “semi-mean” $SMEAN(F) = \int_{-\infty}^{\mu} (x - \mu) dF(x)$.

Lower Partial Moment (LPM)

There are two versions of LPM available, the lower partial moment of order one $LPM1_c$, and the lower partial moment of order two $LPM2_c$, whose functional representation are special cases of the order n lower partial moment

$$LPMn_c(F) = \int_{-\infty}^c (c - x)^n dF(x) \quad (30)$$

where c is a user specified constant “threshold”. This threshold is often referred to as the *minimum acceptable return*, (MAR).

The influence function formulas of these two lower partial moments are:

$$IF(r; LPM1_c; F) = (c - r)\mathbf{I}(r \leq c) - LPM1_c \quad (31)$$

where $LPM1_c = LPM1_c(F)$, and

$$IF(r; LPM2_c; F) = (c - r)^2 \mathbf{I}(r \leq c) - LPM2_c \quad (32)$$

where $LPM2_c = LPM2_c(F)$,

Value-at-Risk (VaR)

The functional form for value-at-risk is

$$VaR_\alpha(F) = -q_\alpha(F). \quad (33)$$

The formula for the influence function of VaR is:

$$IF(r; VaR_\alpha; F) = \frac{1}{f(q_\alpha)} (\mathbf{I}(r \leq q_\alpha) - \alpha) \quad (34)$$

Expected Shortfall (ES)

The functional form of expected shortfall is

$$ES_\alpha(F) = -\frac{1}{\alpha} \int_{-\infty}^{q_\alpha(F)} x \cdot dF(x). \quad (35)$$

The formula for the influence function of the ES is:

$$IF(r; ES; F) = -\frac{I(r \leq q_\alpha)}{\alpha} \cdot (r - q_\alpha) - q_\alpha - ES_\alpha \quad (36)$$

Sharpe Ratio (SR)

The functional representation of the Sharpe ratio is

$$SR(F) = \frac{\mu(F) - r_f}{\sigma(F)} = \frac{\mu_e(F)}{\sigma(F)}. \quad (37)$$

and the influence function for the Sharpe ratio is

$$IF(r; SR; F) = -\frac{\mu_e}{2\sigma^3} (r - \mu)^2 + \frac{1}{\sigma} (r - \mu) + \frac{\mu_e}{2\sigma}$$

where $\mu_e = \mu_e(F) = \mu(F) - r_f$, and $\sigma = \sigma(F)$.

Downside Sharpe Ratio (DSR)

The functional representation of the downside Sharpe ratio is

$$DSR(F) = \frac{\mu(F) - r_f}{\sqrt{2SSD(F)}} = \frac{\mu_e(F)}{\sqrt{2SSD(F)}} \quad (38)$$

and the influence function for $\sqrt{2}DSR(F)$ is

$$IF(r; \sqrt{2}DSR; F) = -\frac{DSR \cdot I(r \leq \mu)}{2 \cdot SemiSD^2} (r - \mu)^2 + \left(\frac{DSR \cdot SemiMean}{SemiSD^2} + \frac{1}{SemiSD} \right) (r - \mu) + \frac{DSR}{2}$$

where $\mu = \mu(F)$, $SemiMean = SemiMean(F)$, $SemiSD = SemiSD(F)$, and $DSR = DSR(F)$. So the influence function of $DSR(F)$ is given by the above expression divided by $\sqrt{2}$.

Sortino Ratio (SoR)

There are two versions of the Sortino ratio. The first version of SoR has the functional representation

$$SoR_c(F) = \frac{\mu(F) - r_f}{\sqrt{LPM2_c(F)}} = \frac{\mu_e(F)}{\sqrt{LPM2_c(F)}} \quad (39)$$

where the subscript c reflects the use of the constant threshold in the denominator lower partial moment of order two.

The second version of SoR has the functional form

$$SoR_\mu(F) = \frac{\mu(F) - r_f}{SSD(F)} = \frac{\mu_e(F)}{SSD(F)} \quad (40)$$

where the subscript μ indicates the use of the mean in the denominator semi-standard deviation.

The influence function for $SoR_c(F)$ is

$$IF(r; SoR_c; F) = \frac{-SoR_c \cdot I(r \leq c)}{2 \cdot LPM2_c} (r - c)^2 + \frac{1}{\sqrt{LPM2_c}} (r - \mu) + \frac{SoR_c}{2} \quad (41)$$

where $\mu = \mu(F)$, $LPM2_c = LPM2_c(F)$, $SoR_c = SoR_c(F)$.

The influence function for $SoR_\mu(F)$ is

$$IF(r; SoR_\mu; F) = -\frac{SoR_\mu \cdot I(r \leq \mu)}{2 \cdot SSD^2} (r - \mu)^2 + \left(\frac{SoR_\mu \cdot SMEAN}{SSD^2} + \frac{1}{SSD} \right) (r - \mu) + \frac{SoR_\mu}{2} \quad (42)$$

where $\mu = \mu(F)$, $SSD = SSD(F)$, $SoR_\mu = SoR_\mu(F)$.

Expected Shortfall Ratio (ESratio)

The functional form is

$$ESratio(F) = \frac{\mu(F) - r_f}{ES_\alpha(F)} = \frac{\mu_e(F)}{ES_\alpha(F)} \quad (43)$$

The formula for the influence function of the ESratio is:

$$IF(r; ESratio; F) = \frac{r - \mu}{ES_\alpha} - \frac{ESratio}{ES_\alpha} \left(-q_\alpha - ES_\alpha - \frac{I(r \leq q_\alpha)}{\alpha} \cdot (r - q_\alpha) \right) \quad (44)$$

Value-at-Risk Ratio (VaRratio)

The functional form of the VaR ratio is:

$$VaRratio(F) = \frac{\mu(F) - r_f}{VaR(F)} = \frac{\mu_e(F)}{-q_\alpha(F)} \quad (45)$$

The formula for the influence function of the VaRratio is:

$$IF(r; VaRratio; F) = -\frac{r - \mu}{q_\alpha} + \frac{VaRratio}{q_\alpha} \times \frac{I(r \leq q_\alpha) - \alpha}{f(q_\alpha)}. \quad (46)$$

Rachev Ratio (RachR)

The functional form is:

$$RachR(F) = \frac{EG_\beta}{ES_\alpha} = \frac{\frac{1}{\beta} \cdot \int_{q_{1-\beta}}^{+\infty} x dF(x)}{-\frac{1}{\alpha} \cdot \int_{-\infty}^{q_\alpha} x dF(x)}$$

where $ES_\alpha = ES_\alpha(F)$ is the expected shortfall at level α and $EG_\beta = EG_\beta(F)$ is the expected tail gain at upper β -quantile defined by the following equation

$$EG_\beta = \frac{1}{\beta} \cdot \int_{q_{1-\beta}}^{+\infty} x dF(x)$$

The formula for the influence function of the Rachev ratio is:

$$IF(r; RaR; F) = \frac{1}{ES_\alpha} \left(\frac{I(r \geq q_{1-\beta})}{\beta} (r - q_{1-\beta}) + q_{1-\beta} - EG_\beta \right) - \frac{RachR}{ES_\alpha} \cdot \left(\frac{-I(r \leq q_\alpha)}{\alpha} (r - q_\alpha) - q_\alpha - ES_\alpha \right) \quad (47)$$

Omega Ratio (Omega)

The functional is:

$$\Omega(F) = \frac{UPM1_c(F)}{LPM1_c(F)} = \frac{\int_c^{+\infty} (x - c) f(x) dx}{\int_{-\infty}^c (c - x) f(x) dx}$$

The formula for the influence function of the Omega ratio is:

$$IF(r; \Omega; F) = \frac{1}{LPM1_c} \cdot \left((r - c) \cdot I(r \geq c) - UPM1_c \right) - \frac{\Omega}{LPM1_c} \cdot \left((c - r) \cdot I(r \leq c) - LPM1_c \right) \quad (48)$$

Appendix B. IF Nuisance Parameters Calculation for Normally Distributed Returns

In order to compute the values of any of the influence functions presented in Section 3, values of the nuisance parameters listed in the third column of Table (2) are needed. One way of obtaining the nuisance parameters is to specify them with “typical” values based an assumed returns distribution. For the risk measure estimators and performance measure estimators, we shall do so here for an assumed returns normal distribution, with monthly mean return of 1% and risk-free rate $r_f = 0$, and monthly volatility of 5% (the corresponding annual mean and volatility are 12% and 17.3%, respectively). We also note that we use $\alpha = 0.10$ for quantiles, VaR and ES, use that α value along with the value $\beta = 0.10$ for the Rachev ratio, and use $c = 0$ for lower partial moments and Sortino ratio with fixed threshold. Below we display the calculation of the corresponding values shown in the third column of Table (2).

SemiSD

$\sigma^2 = 2 \cdot \sigma_{semisd}^2 = 2 \cdot SemiSD^2$, and so $SSD = \sigma/\sqrt{2}$. Thus $SemiSD = 0.0353$.

SMean

$SMean = \int_{-\infty}^{\mu} (r - \mu) f(r) dr$, and for $f(r|\mu, \sigma) = \frac{1}{\sqrt{2\pi}} \cdot \exp\left(-\frac{(r - \mu)^2}{2\sigma^2}\right)$, we have $SMean = \frac{-\sigma}{\sqrt{2\pi}} = -0.0199$.

LPM1

Under normal distribution $N(\mu, \sigma^2)$, $LPM1_c = LPM1_c(\mu, \sigma)$ as a function of μ and σ is

$$\begin{aligned} LPM1_c(\mu, \sigma) &= \int_{-\infty}^c (c - x) f(x|\mu, \sigma) dx \\ &= \int_{-\infty}^d (d - y) f(y|0, 1) dy \cdot \sigma \\ &= (d \cdot \Phi(d) + \phi(d)) \cdot \sigma \end{aligned}$$

where $d = \frac{c - \mu}{\sigma}$ and Φ and ϕ are the cdf and pdf of the standard normal distribution function.

For the given values of μ , σ , and $c = 0$, we have:

$$LPM1_c = .0153.$$

LPM2

For the normal distribution $N(\mu, \sigma^2)$, $LPM2_c = LPM2_c(\mu, \sigma)$ as a function of μ and σ is

$$\begin{aligned} LPM2_c(\mu, \sigma) &= \int_{-\infty}^c (c - x)^2 f(x|\mu, \sigma) dx \\ &= \int_{-\infty}^d (d - y)^2 f(y|0, 1) dy \cdot \sigma^2 \\ &= ((d^2 + 1) \cdot \Phi(d) + d \cdot \phi(d)) \cdot \sigma^2 \end{aligned}$$

where $d = \frac{c - \mu}{\sigma}$ and Φ and ϕ are the cdf and pdf of the standard normal distribution function. For the given value of $\mu = 1\%$, $\sigma = 5\%$ and $c = 0$, we have

$$LPM2_c = .000898.$$

Expected Shortfall

The formula for expected shortfall for a normal distribution is

$$ES_\alpha = -\mu + \sigma \cdot \frac{\phi(z_\alpha)}{\alpha}$$

where ϕ is the standard normal density function and z_α is the lower α -quantile of the standard normal distribution. For the specified parameters, we have $ES_\alpha = .0777$.

Alpha Quantile

For the specified normal distribution, this quantile is $q_\alpha = q_{.10} = -0.0541$.

VaR

For VaR the nuisance parameter $f(q_\alpha)$ is the normal density $f_{normal}(r; \mu, \sigma^2)$ evaluated at $r = q_\alpha$, and for the specified parameters we have $f_{normal}(q_\alpha; \mu, \sigma^2) = 3.51$.

Sharpe Ratio (SR)

For the specified normal distribution with $\mu = \mu_e = 1\%$ and $\sigma = 5\%$, the Sharpe ratio is $SR = 0.2$.

Downside Sharpe Ratio (DSR)

For the specified normal distribution with $\mu = \mu_e = 1\%$ and $\sigma = 5\%$, we have $SemiSD = \sigma = .05$, $SMEAN = -\phi(0) \cdot \sigma = -.0199$, and $DSR = \frac{\mu_e}{\sqrt{2}SemiSD} = .2$.

Sortino Ratio (SoR) with Mean Threshold

For the specified normal distribution with $\mu = \mu_e = 1\%$ and $\sigma = 5\%$, we have $SemiSD = \sigma/\sqrt{2} = .0354$, $SMEAN = -\phi(0) \cdot \sigma = -.0199$, and $SoR_\mu = \frac{\mu_e}{SemiSD} = .2828$.

Sortino Ratio (SoR) with Constant Threshold c

For the specified normal distribution with $\mu = \mu_e = 1\%$ and $\sigma = 5\%$, $d = \frac{c - \mu}{\sigma} = -.2$. Therefore $LPM2_c = ((d^2 + 1) \cdot \Phi(d) + d \cdot \phi(d)) \cdot \sigma^2 = .000898$.

Sortino ratio with constant threshold c is $SoR_c = \frac{\mu_e}{\sqrt{LPM2_c}} = .3337$.

Expected Shortfall Ratio (ESratio)

For the specified normal distribution with $\mu = \mu_e = 1\%$ and $\sigma = 5\%$, we have $q_\alpha = q_{.10} = -.0541$. The expected shortfall is $ES_\alpha = -\mu + \sigma \cdot \frac{\phi(z_\alpha)}{\alpha} = .0777$ and ES ratio is $ESratio = \frac{\mu_e}{ES_\alpha} = .129$.

VaR Ratio (VaRratio)

For the specified normal distribution with $\mu = \mu_e = 1\%$ and $\sigma = 5\%$, we have $q_\alpha = q_{.10} = -.0541$. The VaR ratio is $VaRratio = \frac{\mu_e}{-q_\alpha} = .185$.

Rachev Ratio (RachR)

For the normal distribution $N(\mu, \sigma^2)$, the upper β -quantile is $q_{1-\beta} = \mu + z_{1-\beta} \cdot \sigma = .01 + 1.2816 \times .05 = .0741$. The expected tail gain at upper β -quantile is

$$\begin{aligned}
 EG_\beta &= \frac{1}{\beta} \cdot \int_{q_{1-\beta}}^{+\infty} x f(x|\mu, \sigma) dF(x) \\
 &= \frac{1}{\beta} \cdot \int_{z_{1-\beta}}^{+\infty} (\mu + \sigma y) f(y|0, 1) dy \\
 &= \frac{\mu}{\beta} \cdot \int_{z_{1-\beta}}^{+\infty} f(y|0, 1) dy + \frac{\sigma}{\beta} \cdot \int_{z_{1-\beta}}^{+\infty} y \cdot f(y|0, 1) dy \\
 &= \mu + \frac{\sigma}{\beta} \cdot \int_{z_{1-\beta}}^{+\infty} y \cdot \frac{1}{\sqrt{2\pi}} \cdot \exp\left(\frac{-y^2}{2}\right) dy \\
 &= \mu + \frac{\sigma}{\beta} \cdot \frac{1}{\sqrt{2\pi}} \cdot \exp\left(\frac{-(z_{1-\beta})^2}{2}\right) \\
 &= \mu + \frac{\sigma}{\beta} \cdot \phi(z_{1-\beta})
 \end{aligned}$$

With the specified parameters, $EG_\beta = .0977$.

The Rachev ratio is $RachR = \frac{EG_\beta}{ES_\alpha} = \frac{.0977}{.0777} = 1.257$.

Omega Ratio (Omega)

For the given parameters, the LPM1 is $LPM1_c(\mu, \sigma) = (d \cdot \Phi(d) + \phi(d)) \cdot \sigma = .0153$.

The upper partial moment is

$$\begin{aligned}UPM1_c(\mu, \sigma) &= \int_c^{+\infty} (x - c) f(x|\mu, \sigma) dx \\&= \int_{-\infty}^{+\infty} (x - c) f(x|\mu, \sigma) dx - \int_{-\infty}^c (x - c) f(x|\mu, \sigma) dx \\&= \mu - c + \int_{-\infty}^c (c - x) f(x|\mu, \sigma) dx \\&= \mu - c + LPM1_c(\mu, \sigma)\end{aligned}$$

Therefore $UPM1_c(\mu, \sigma) = .01 - 0 + .0153 = .0253$ and the Omega ratio is $\Omega = \frac{.0153}{.0253} = 1.652$.

Code for Above Calculations

The following code is in essence used in the RPEIF package function `nuisParsFn`.

```
#-----  
# Nuisance Parameters - Risk Estimators  
#-----  
  
# Mean  
mu <- 0.01  
sd <- 0.05  
c <- 0  
alpha <- 0.1  
  
# SemiSD  
SemiSD <- sd/sqrt(2)  
SemiSD  
  
## [1] 0.03535534  
  
SMEAN <- -dnorm(0) * sd  
SMEAN  
  
## [1] -0.01994711
```

```

# LPM1
d <- (c - mu)/sd
LPM1 <- (d * pnorm(d) + dnorm(d)) * sd
LPM1

## [1] 0.01534473

# LPM2
d <- (c - mu)/sd
LPM2 <- ((d^2 + 1) * pnorm(d) + d * dnorm(d)) * sd^2
LPM2

## [1] 0.0008984034

# ES
q_alpha <- mu + qnorm(alpha) * sd
q_alpha

## [1] -0.05407758

ES <- -mu + dnorm(qnorm(alpha))/alpha * sd
ES

## [1] 0.07774917

# VaR
q_alpha <- mu + qnorm(alpha) * sd
q_alpha

## [1] -0.05407758

f_q_alpha <- dnorm(q_alpha, mu, sd)
f_q_alpha

## [1] 3.509967

#-----
# Nuisance Parameters - Performance Estimators
#-----

# Mean
mu <- 0.01
sd <- 0.05

```

```

c <- 0
alpha <- 0.1
beta <- 0.1

# SR
SR <- mu/sd
SR

## [1] 0.2

# SoR_c
d <- (c - mu)/sd
LPM2 <- ((d^2 + 1) * pnorm(d) + d * dnorm(d)) * sd^2
LPM2

## [1] 0.0008984034

SoRc <- mu/sqrt(LPM2)
SoRc

## [1] 0.3336294

# SoR_mu
SemiSD <- sd/sqrt(2)
SemiSD

## [1] 0.03535534

SMEAN <- -dnorm(0) * sd
SMEAN

## [1] -0.01994711

SoRmu <- mu/SemiSD
SoRmu

## [1] 0.2828427

# ESratio
q_alpha <- mu + qnorm(alpha) * sd
q_alpha

## [1] -0.05407758

```

```

ES <- -mu + dnorm(qnorm(alpha))/alpha * sd
ES

## [1] 0.07774917

ESratio <- mu/ES
ESratio

## [1] 0.1286187

# VaRatio
q_alpha <- mu + qnorm(alpha) * sd
q_alpha

## [1] -0.05407758

VaRatio <- -mu/q_alpha
VaRatio

## [1] 0.1849195

f_q_alpha <- dnorm(q_alpha, mu, sd)
f_q_alpha

## [1] 3.509967

# RachevRatio
q_alpha <- mu + qnorm(alpha) * sd
q_alpha

## [1] -0.05407758

ES <- -mu + dnorm(qnorm(alpha))/alpha * sd
ES

## [1] 0.07774917

q_one_minus_beta <- mu + qnorm(1 - beta) * sd
q_one_minus_beta

## [1] 0.07407758

```



```

EG <- mu + dnorm(qnorm(1 - beta))/beta * sd
EG

## [1] 0.09774917

RachR <- EG/ES
RachR

## [1] 1.257237

# OmegaRatio
d <- (c - mu)/sd
LPM1 <- (d * pnorm(d) + dnorm(d)) * sd
LPM1

## [1] 0.01534473

UPM1 <- LPM1 + mu - c
UPM1

## [1] 0.02534473

Omega <- UPM1/LPM1
Omega

## [1] 1.651689

```

References

- X. Chen and R. D. Martin. Standard errors of risk and performance measure estimators for serially correlated returns. *Journal of Risk*, 23:1–41, 2021.
- F R Hampel. The influence curve and its role in robust estimation. *Journal of the American Statistical Association*, 69(346):383–393, 1974.
- F R Hampel, E M Ronchetti, P J Rousseeuw, and W A Stahel. *Robust Statistics: The Approach Based on Influence Functions*. Wiley, 1986.
- RA Maronna, RD Martin, VJ Yohai, and M Salibian-Barrera. Robust statistics: Theory and methods (with r) wiley. *Hoboken, NJ, USA*, 2019.
- R D Martin and S Zhang. Nonparametric versus parametric expected shortfall. *Journal of Risk*, 21(6):1–41, 2019. URL <https://ssrn.com/abstract=2747179>.

- S. Zhang. Statistical analysis of portfolio risk and performance measures: the influence function approach. 2009.
- S. Zhang, R. D. Martin, and A. A. Christidis. Influence functions for risk and performance estimators. 2020.
URL <https://ssrn.com/abstract=3415903>.